

WHAT IS CLAIMED IS:

1. A method for identifying an unknown base sequence present in a target single-stranded nucleic acid comprising the steps of:

5 (a) preparing a probe array in which single-stranded nucleic acid probes of No. 1 to No. n ( $n \geq 2$ ) are arranged as isolated spots on a substrate, the probes each having a base sequence complementary to one of plural base sequences expected to be the unknown  
10 base sequence;

(b) reacting a single-stranded nucleic acid, which has a base sequence fully complementary to a base sequence of one of the single-stranded nucleic acid probes and is fluorescence-labeled, with the probe  
15 array under such conditions that single-stranded nucleic acids complementary to each other form a double-stranded nucleic acid;

removing the unreacted labeled single-stranded nucleic acid, and  
20 measuring fluorescence intensity of each spot of the probe array to obtain a first template pattern showing a relationship between location of the probes and fluorescent characteristics;

(c) performing the same operation as the step (b)  
25 for each of remaining single-stranded nucleic acid probes using a second to a nth single-stranded nucleic acid, and obtaining template patterns of No. 2 to No. n

004495-0340  
1000-953400

showing a relationship between location and fluorescent characteristics of the probes;

(d) performing the same operation as the step (b) using a sample containing the target single-stranded nucleic acid of unknown base sequence to obtain a sample pattern showing relationship between a position and fluorescent characteristics; and

(e) comparing the sample pattern obtained in the step (d) with n pieces of template patterns obtained in the steps (b) and (c), to identify a template pattern showing substantially the same pattern as the sample pattern and identifying the base sequence of the single-stranded nucleic acid used for the preparation of the identified template pattern as the unknown base sequence of the target single-stranded nucleic acid.

2. A method for identifying an unknown base sequence present in a target single-stranded nucleic acid comprising the steps of:

(a) preparing a probe array in which single-stranded nucleic acid probes of No. 1 to No. n ( $n \geq 2$ ) are arranged as isolated spots on a substrate, the probes each having a base sequence complementary to one of plural base sequences expected to be the unknown base sequence;

(b) reacting a single-stranded nucleic acid which has a base sequence fully complementary to a base

sequence of one of the single-stranded nucleic acid probes and is fluorescence-labeled, with the probe array under such conditions that single-stranded nucleic acids complementary to each other form a double-stranded nucleic acid;

removing the unreacted labeled single-stranded nucleic acid, and

measuring fluorescence intensity of each spot of the probe array to obtain a first template pattern showing a relationship between location of the probes and fluorescent characteristics;

(c) analyzing the first template pattern to locate probes and to calculate a mean value of fluorescence intensities ( $F_i$ ) of the double-stranded nucleic acids having  $i$  of mismatched base pairs, where  $i$  is an integer not less than 1;

(d) calculating a difference ( $F_1, 0$ ) between the fluorescence intensity of the fully complementary double-stranded nucleic acid without mismatch ( $F_0$ ) and the mean value of the fluorescence intensities of the double-stranded nucleic acids having one-base mismatch ( $F_1$ ), further calculating a difference ( $F_{i+1}, i$ ) between a fluorescence intensity of a double-stranded nucleic acid having  $(i+1)$  base mismatches ( $F_{i+1}$ ) and a fluorescence intensity of a double-stranded nucleic acid having  $i$ -base mismatches ( $F_i$ ), and identifying  $i$  being  $F_{i+1}, i < F_i, i-1$ ;

0042596-083104  
TOP SECRET

(e) assuming a target DNA which base sequence is complementary to the second probe sequence, then obtaining the second template pattern formed by the probe position where the number of mismatched base pairs to the target having the complementary sequence to the second probe sequence is not more than  $i$ ;

(f) performing the same operation as the step (e) for each of remaining single-stranded nucleic acid probes using a third to a  $n$ th single-stranded nucleic acid, and obtaining template patterns of No. 3 to No.  $n$  showing a relationship between location and fluorescent characteristics of the probes, wherein the template patterns are formed from the positions of the probes having a base sequence that forms mismatched base pairs in a number not more than  $i$ ;

(g) performing the same operation as the step (b) using a sample containing the target single-stranded nucleic acid of unknown base sequence to obtain a sample pattern showing relationship between a position and fluorescent characteristics; and

(h) comparing the sample pattern obtained in the step (g) with  $n$  pieces of template patterns obtained in the steps (b), (c) and (e), to identify a template pattern showing essentially the same pattern as the sample pattern and identifying the base sequence of the single-stranded nucleic acid used for the preparation of the identified template pattern as the unknown base

00412506 083104  
FOR EEO 92501650

sequence of the target single-stranded nucleic acid.

3. The method according to claim 2, wherein the  
step (g) further comprises the substep of obtaining a  
5 two-valued pattern of the fluorescence intensity by  
using the threshold fluorescence intensity  $F_i$ .

4. The method according to claim 2, wherein the  
length of the probe is 8 mer to 30 mer.

5. The method according to claim 4, wherein the  
length of the probe is 12 mer to 25 mer.

6. The method according to claim 2, wherein the  
15 number of the mismatched base pairs (i) is 1.

7. The method according to claim 1, wherein in  
the steps (b), (c) and (d), the probes in the probe  
array are heat-denatured in a solution containing the  
20 single-stranded nucleic acid, and cooled to a  
temperature suitable for a double-stranded formation  
reaction while the substrate is soaked in the solution.

8. The method according to claim 7, wherein the  
25 length of the single-stranded nucleic acid probe is 18  
mer, the temperature for performing the heat  
denaturation is 70°C or more, the temperature for the

004456-003401

double-strand formation reaction is 40°C or more, and  
100 mM sodium chloride is contained in the sample  
solution at that time.

000121506 003401  
T07E00 90527600